

Searching the charged Higgs boson of the type III two Higgs doublet model

H. Cardenas^{1,2} and J-Alexis Rodríguez²

¹*Escuela de Física, Universidad Pedagógica y Tecnológica de Tunja, Duitama, Colombia*

²*Departamento de Física, Universidad Nacional de Colombia
Bogotá, Colombia*

In the framework of the Two Higgs Doublet Model (2HDM) type III appears two charged Higgs boson and recently there are experimental reports from D0 and CDF collaborations searching a particular signature of new physics. We present a review of the analysis done in the region $M_{H^+} > m_t$ by D0 collaboration and we use the ratio R_σ for the region $M_{H^+} < m_t$ in different scenarios of space parameter of this model.

The Two Higgs Doublet Model (2HDM) is a simple extension of the standard model (SM), where an additional Higgs doublet is introduced [1]. The two complex scalar fields correspond to eight degrees of freedom, three of them are absorbed as Goldstone bosons in order to give mass and as longitudinal components to the $W^\pm$ and Z bosons. The remaining degrees of freedom are interpreted as five physical states: two neutral scalars h^0 and H^0 , a pseudo-scalar A^0 , and a pair of charged Higgs bosons $H^\pm$. Another consequence of this extension is a more generic pattern of Flavor Changing Neutral Currents (FCNC), including such processes at tree level. Because of low energy experiments these FCNC could be a problem, usually solved by imposing a discrete symmetry that allows to couple at most one Higgs doublet to each fermion. In the 2HDM called type I, one Higgs doublet gives masses to the up and down quarks, simultaneously. And, in the model type II, one Higgs doublet gives masses to the up type quarks and the other one to the down type quarks, and it is precisely the structure of the minimal supersymmetric extension of the SM (MSSM). However, the addition of these discrete symmetries is not mandatory and in that case both doublets contribute to generate the masses for up-type and down-type quarks, such model is known as the 2HDM type III [2].

While it may be hard to distinguish any one of these neutral Higgs bosons of the 2HDM-III from that one of the SM, the charged $H^\pm$ pair carry a distinctive hall-mark of this kind of new physics. Hence the charged Higgs boson plays a very important role in the search of new physics beyond the SM.

Usually the limits on the charged Higgs mass are in the context of the type II 2HDM. Direct searches have carried out by LEP experiments, searching for pair produced charged Higgs bosons via s -channel exchange of a Z -boson or a photon. LEP collaborations yield a combined lower limit on $M_{H^\pm}$ of 78.6 GeV [3] assuming $H^+ \rightarrow \tau^+ \nu_\tau (c\bar{s})$ in a wide range of the ratio of the vacuum expectation values of the two Higgs doublets, $\tan\beta = v_2/v_1$. Searches in the hadron colliders consider two cases $M_{H^\pm} > m_t$ and $M_{H^\pm} < m_t$. At the Tevatron, the direct searches for charged Higgs boson are based on $p\bar{p} \rightarrow t\bar{t}$ with at least one top quark using $t \rightarrow H^+ b$, for the first case $M_{H^\pm} > m_t$. The CDF collaboration has reported a direct search for charged Higgs boson decaying into $\tau^+ \nu_\tau$, $c\bar{s} t\bar{b}$ or $W^+ A^0$, which uses measurements of the top pair production cross section in the $leptons + \cancel{E}_T + jets + leptons$ channels, from data samples corresponding to an integrated luminosity of 193 pb⁻¹ [4]. On the other hand, recently the D0 collaboration has presented a direct search for a charged Higgs boson produced by $q\bar{q}$ annihilation and decaying to $t\bar{b}$ final state, in the $180 \leq M_{H^+} \leq 300$ GeV mass range, it is $m_{H^+} > m_t$, using 0.93 fb⁻¹ of data collected at center-of-mass energy $\sqrt{s} = 1.96$ TeV [5]. The analysis lead to upper limits on the production cross section in the 2HDM types I, II and III [5, 6]. We are going to discuss different scenarios for the 2HDM type III using these experimental results. Also, searches in the $M_{H^\pm} < m_t$ region have been performed, using the production cross section of top quark pairs at the Tevatron [6, 7], and we intend to exploit this fact to discuss the possible limits on the charged Higgs boson mass obtained from measurements of the ratio $R_\sigma = \sigma_{t\bar{t}}^{l+jets} / \sigma_{t\bar{t}}^{dileptons}$ in the framework of the 2HDM type III.

Other experimental bounds on the charged Higgs boson mass come from virtual charged Higgs boson production in $b \rightarrow s\gamma$ transitions [8]. Finally, the search for the charged Higgs boson will continue above the top quark mass at the experiments of LHC, ATLAS and CMS. The main production mechanisms would be the processes $gg \rightarrow tbH^+$ and $gb \rightarrow tH^+$ which have been studied using simulations of the LHC detectors [9].

In the 2HDM both doublets can acquire vacuum expectation value (VEV) different from zero. As mentioned before, we will focus our study to the 2HDM type-III, which owns the most general Yukawa Lagrangian (without additional symmetries imposed), given by [10, 11, 12]

$$-\mathcal{L}_Y^{(III)} = \eta_{ij}^{U,0} \bar{Q}_{iL}^0 \tilde{\Phi}_1 U_{jR}^0 + \eta_{ij}^{D,0} \bar{Q}_{iL}^0 \Phi_1 D_{jR}^0 + \xi_{ij}^{U,0} \bar{Q}_{iL}^0 \tilde{\Phi}_2 U_{jR}^0 + \xi_{ij}^{D,0} \bar{Q}_{iL}^0 \Phi_2 D_{jR}^0 + \text{h.c.}, \quad (1)$$

where Φ_i are the complex scalar fields, η_{ij}^0 and ξ_{ij}^0 are complex (even if later on we simplify the problem to real λ , the original coupling matrices can be also complex as in the Yukawa sector of the SM) and non-diagonal coupling matrices and the suffix "0" denotes flavor eigenstates.

The scalar spectrum has mass eigenstates which contain two CP -even neutral Higgs bosons (h^0, H^0) coming from the mixing of $\Re(\phi_i^0)$ with mixing angle α ; two charged Higgs bosons ($H^\pm$) which corresponds to a mixture of the

would-be Goldstone bosons $G_W^\pm$ through the mixing angle $\tan\beta = v_2/v_1$; and one CP -odd Higgs (A^0) which mix the neutral would-be Goldstone boson G_Z^0 . In the framework of the type III 2HDM there is a global symmetry which can make a rotation of the Higgs doublets and fix one VEV equal to zero [10]. In such a way, $v_1 = v$ and $v_2 = 0$, and the mixing parameter $\tan\beta = v_2/v_1$ can be eliminated from the Lagrangian. It is worth to mention that the experimental sensitivities are usually displayed in terms of the parameters m_A and $\tan\beta$ or $M_{H^\pm}$ and $\tan\beta$, that characterize the MSSM Higgs sector or the 2HDM type II. If the parameter $\tan\beta$ is eliminated from the Lagrangian, we have the usual 2HDM type III [10], where the Lagrangian of the charged sector is given by

$$-\mathcal{L}_{H^\pm ud}^{III} = H^+ \bar{U} [K \xi^D P_R - \xi^U K P_L] D + h.c. \quad (2)$$

where K is the Cabbibo-Kobayashi-Maskawa (CKM) matrix and $\xi^{U,D}$ the flavor changing matrices. In the present work we take into account the Cheng-Sher-Yuan (CSY) parameterization which is built as the geometrical mean of the Yukawa couplings of the quark fields, $\xi_{ij} \equiv \frac{\sqrt{m_i m_j}}{v} \lambda_{ij}$ [11].

The search of D0 collaboration [5] use the process $q\bar{q}' \rightarrow H^+ \rightarrow t\bar{b}$ followed by $t \rightarrow b(W^+ \rightarrow l\nu)$, which correspond to a single top production, with a final signature 2- b -tagged +lepton+ missing energy. We notice there are two vertices type $H^+ q\bar{q}'$ where appear the parameters $\xi_{qq'}$, and then different considerations can be done, taking into account that appears terms like $\sum_j (K_{qj} \xi_{jq'})$. In the annihilation vertex $q\bar{q}' H^+$, it is possible in the framework of the 2HDM type III: $c\bar{b}H^+$ and $c\bar{s}H^+$; about this point H. He and C. P. Yuan [13] showed that it is possible to enhanced the production cross section if the vertex $c\bar{b}H^+$ is important, and it is possible if the parameter λ_{tc} is bigger than one. We have evaluated the option of the vertex $c\bar{s}H^+$ in the framework of the 2HDM-III and it is smaller two or three orders of magnitude than the vertex $c\bar{b}H^+$. Now in the second vertex, the H^+ decay vertex, we have several couplings regarding the 2HDM-III allowed processes: $H^+ \rightarrow t\bar{s}$, $H^+ \rightarrow c\bar{b}$ and $H^+ \rightarrow t\bar{b}$. The first option, into $t\bar{s}$, does not have a quark b in the final state, and the experiment is looking for at least two b -tagged, one coming from the H^+ decay and the second one coming from the top quark decay. The second option is into $c\bar{b}$ but it is doubly suppressed for the factor ξ_{tc} and for the Cabbibo-Kobayashi-Maskawa K_{cb} factor. Finally, we will have the same main channel production in the 2HDM type III that in the types I and II, $c\bar{c} \rightarrow H^+ \rightarrow t\bar{b}$ followed by $t \rightarrow (W^+ \rightarrow l\nu)b$.

Using the parameterization CYS [11] and the numerical values of the CKM matrix K_{ij} , the flavor changing couplings, $\sum_j K_{qj} \xi_{jq'}$, of the charged Higgs boson to the fermions are shown in Tab. I. These factors have been simplified taken into account the numerical values of quark masses and assuming that the parameters λ_{ij} involving the first and second generations are smaller than parameters involving the third generation. Finally, bounds and restrictions on the λ_{ij} for the quark sector and ξ_{ij} for the leptonic sector can be found in literature [10, 14].

The analysis presented by D0 collaboration [5] for the type III 2HDM has followed the analysis done in reference [13] assuming that the parton level is important to enhance the cross section if λ_{tc} is bigger than one, the experimental analysis have used $\lambda_{tc} = 5$ [6]. Further, they assumed the parameter λ_{tt} in the charged Higgs decay vertex, to be equal to λ_{tc} . About this last point, we should mention that Atwood, *et. al.* in reference [10] have already shown that the assumption $\lambda_{ij} = \lambda$ is not in agreement with the low energy phenomenology and on the other hand, it has been shown [15] that perturbation theory validity requires that $\lambda_{tc} \leq 2.8$. From this point we aim to explore scenarios




FIG. 1: The cross section times branching fraction versus the charged Higgs boson for different scenarios in the type III 2HDM

(ij)	$(\xi^U K)_{ij}$	$(K\xi^D)_{ij}$
$c\bar{s}$	$K_{ts} \frac{\lambda_{ct}\sqrt{m_c m_t}}{v}$	
$c\bar{b}$	$K_{tt} \frac{\lambda_{ct}\sqrt{m_c m_t}}{v}$	$K_{cb} \frac{\lambda_{bb} m_b}{v}$
$t\bar{s}$	$K_{ts} \frac{\lambda_{tt} m_t}{v}$	
$t\bar{b}$	$K_{tb} \frac{\lambda_{tt} m_t}{v}$	$K_{tb} \frac{\lambda_{bb} m_b}{v}$

TABLE I: The couplings of the vertices $H^+ q\bar{q}'$ are proportional to $\sum_j (K_{qj} \xi_{jq'})$ but they can be simplified taking into account the CSY parameterization and the numerical values of the K_{CKM} matrix as it is shown.

allowed in the 2HDM-III space parameters, with the additional simplification that in this model $\tan\beta$ is a spurious parameter. The experimental D0 collaboration report the observed limits on the production cross section (pb) times the branching fraction, $\sigma(q\bar{q}' \rightarrow H^+) \times B(H^+ \rightarrow t\bar{b})$, these limits are shown in Fig. 1. We have used the program CompHEP [16] to simulate charged Higgs boson production and decays, $q\bar{q}' \rightarrow H^+ \rightarrow t\bar{b} \rightarrow W^+ b\bar{b} \rightarrow l^+ \nu b\bar{b}$ where l represents an electron or muon. The expected limits using the same values of the λ_{ij} in the charged Higgs boson mass range 180 to 300 GeV are plotted in Fig. 1. In addition, here can be seen the predictions taken $\lambda_{tc} = 2.8, 1$, $\lambda_{tt} = 2.8, 1$ and $\lambda_{bb} = 0, 10$. We can conclude that restrictions on the parameter space of the type III 2HDM are not too stronger. We should notice that in the case $\lambda_{ij} = 5$ only charged Higgs masses above around 264 GeV are allowed and for values $\lambda_{tt} = 2.8$ the charged Higgs boson mass should be bigger than 230 GeV. For completeness the branching fraction ratios in the different scenarios considered are shown in Fig. 2, where we have use the parameters $\xi_{\tau\tau} = 2.2 \times 10^{-2}$, $\xi_{\tau\mu} = 2.1 \times 10^{-1}$, and $\xi_{\mu\mu} = 0.12$ according to the allowed regions cited in the literature [10, 14].

The discussion presented until now is for charged Higgs boson mass bigger than the top quark mass, but we can also ask for the contrary case $M_{H^+} < m_t$. In this case the decay $t \rightarrow H^+ b$ compete with the standard decay channel $t \rightarrow W^+ b$ and the experimental analysis has been done using the cross section of the top quark pair production [6, 7]. In order to estimate upper limits on the branching fraction $B(t \rightarrow H^+ b)$ is useful to use the ratio $R_\sigma = \sigma(p\bar{p} \rightarrow t\bar{t})_{l+jets} / \sigma(p\bar{p} \rightarrow t\bar{t})_{dilepton}$. The ratio R_σ should be consistent with unity if the dilepton and lepton+jets analyses are in the framework of the standard model $t\bar{t}$ production. The ratio R_σ will have smaller systematic uncertainties than




FIG. 2: Branching ratios in the 2HDM-III for the parameters defined in each figure.

individual cross section measurements and also some common factors will cancel. And of course, R_σ is independent of any theory prediction for $\sigma(p\bar{p} \rightarrow t\bar{t})$, so it is a better place to look for new physics by looking for a deviation in the ratio, rather than by comparing a measured cross section to a theory prediction. R_σ is sensitive to decays such as $t \rightarrow H^+b$, it is possible to give an interpretation in terms of the branching fraction and a measurement of R_σ can be translate into an upper limit on $B(t \rightarrow H^+b)$. For instance if we consider that $B(H^+ \rightarrow cs) = 100\%$ then we can get $B(t \rightarrow H^+b) = 0.13 \pm 0.12$ in the type II 2HDM. In general this ratio can be written according to [6, 17]

$$R_\sigma = 1 + \frac{B(t \rightarrow H^+b)}{[1 - B(t \rightarrow H^+b)]B(W^+ \rightarrow q\bar{q}')} . \quad (3)$$

Experimentally this ratio has been measured by D0 collaboration $R_\sigma = 1.21^{+0.27}_{-0.26}$ [6] and by the CDF collaboration $1/R_\sigma = 1.45^{+0.83}_{-0.55}$ [7]. We have plotted the ratio R_σ in figure 3 in the framework of the 2HDM type III for the scenarios that we used so far.




FIG. 3: The ratio R_σ with the experimental limits from D0 collaboration and different values of the space parameter in the framework of the type III 2HDM

As a conclusion we have reviewed the analysis presented by D0 collaboration [5] in the first search for a charged Higgs boson directly produced by quark-antiquark annihilation and decaying into the $t\bar{b}$ final state, in the $180 \leq M_{H^+} \leq 300$ GeV, analyzing 0.9 fb^{-1} of data from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, in the framework of the 2HDM type III. We have explored different scenarios of the space parameter in the region $M_{H^+} > m_t$ and in the region $M_{H^+} < m_t$ we have used the measurements of the ratio R_σ [6, 7]. Results are shown in figures 1 and 3.

We would like to acknowledge to Y. Peters for useful discussions about the experimental results from her talk presented in SUSY08 conference (Seoul, Korea, June 2008) and to C. Jimenez for his valuable help with data manipulation. Also we want to acknowledge to D. Milanes for his careful reading of the manuscript and useful suggestions.

-
- [1] J. Gunion, *et. al.*, *The Higgs Hunter's Guide*, Frontiers in Physics (Addison-Wesley, Redwood City, CA, 1990).
 - [2] S. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977); W. S. Hou, Phys. Lett. B **296**, 179 (1992); D. Chang, W. S. Hou and W. Y. Keung, Phys. Rev. D **48**, 217 (1993) [arXiv:hep-ph/9302267].
 - [3] LEP Higgs Working Group for Higgs boson searches, arXiv:hep-ex/0107031; P. Achard, *et. al.*, [L3 Collaboration], Phys. Lett. B **575**, 208 (2003) [arXiv:hep-ex/0309056]; A. Heister, *et. al.*, [ALEPH Collaboration], Phys. Lett. B **543**, 1 (2002) [arXiv:hep-ex/0207054]; J. Abdallah, *et. al.*, [DELPHI Collaboration], Eur. Phys. J. C **34**, 399 (2004) [arXiv:hep-ex/0404012].
 - [4] A. Abulencia, *et. al* [CDF Collaboration], Phys. Rev. Lett. **96**, 042003 (2006) [arXiv:hep-ex/0510065].
 - [5] V. M. Abazov, *et. al* [D0 Collaboration], arXiv:hep-ex/0807.0859.
 - [6] Y. Peters, Search for charged Higgs boson at D0, Talk presented in SUSY08 conference, 16-21 June 2008, Seoul, Korea, [arXiv:hep-ex/0810.2078].
 - [7] K. Bloom, arXiv:hep-ex/0405020.
 - [8] F. M. Borzumati and C. Greub, Phys. Rev. D **58**, 074004 (1998) [arXiv:hep-ph/9802391]; Z. j. Xiao and L. Guo, Phys. Rev. D **69**, 014002 (2004) [arXiv:hep-ph/0309103].

- [9] M. Carena, S. Heinemeyer, C. Wagner and G. Weiglein, *Eur. J. Phys. C* **45**, 797 (2006) [arXiv:hep-ph/0511023]; K. Assamagan, Y. Coadou and A. Deandrea, *Eur. J. Phys. C* **4**, 9 (2002); K. Assamagan and N. Gollub, *Eur. J. Phys. C* **39S2**, 25 (2005) [arXiv:hep-ph/0406013]; M. Hashemi, S. Heinemeyer, R. Kinnunen, A. Nikitenko and G. Weiglein, arXiv:hep-ph/0804.1228.
- [10] D. Atwood, L. Reina and A. Soni, *Phys. Rev. D* **55**, 3156 (1997) [arXiv:hep-ph/9609279] and references there in.
- [11] T. P. Cheng and M. Sher, *Phys. Rev.* **D35**, 3484 (1987).
- [12] R. A. Diaz, R. Martinez and J-Alexis Rodriguez, *Phys. Rev.* **D63**, 095007 (2001), [arXiv:hep-ph/0010149].
- [13] H. He and C. P. Yuan, *Phys. Rev. Lett.* **83**, 28 (1999).
- [14] R. A. Diaz, R. Martinez and C. E. Sandoval, [arXiv:hep-ph/0311201] (2003); R. A. Diaz, R. Martinez and C. E. Sandoval, [arXiv:hep-ph/0406265] (2004); R. A. Diaz, R. Martinez and J-Alexis Rodriguez, [arXiv:hep-ph/0103050] (2001); R. A. Diaz, R. Martinez and J-Alexis Rodriguez, *Phys. Rev.* **D63**, 095007 (2001), [arXiv:hep-ph/0010149].
- [15] R. Martinez, J. Alexis Rodriguez and M. Roza, *Phys. Rev. D* **68**, 035001 (2003) [arXiv:hep-ph/0212236].
- [16] E. Boos *et al.* [CompHEP Collaboration], *Nucl. Instrum. Meth. A* **534**, 250 (2004) [arXiv:hep-ph/0403113].
- [17] A. Quadt, *Eur. J. Phys. C* **48**, 835 (2006).